

Plantar Fasciitis

The plantar fascia runs along the length of the sole of the foot and along with various ligaments and muscles of the foot, functions to assist in maintaining its inner arch (like the string of a bow, with the bones of the foot being the bow itself).

Plantar fasciitis is an inflammation of the origin of the fascia at the calcaneus (heel bone). It is a result of a repetitive strain / tear and repair of the fascia from traction forces that have worn it out. It can be likened to an elastic band that has been left out in the sun and stretched too many times; it loses its stretch and has many micro-tears in it.

Heel spurs have long been associated with heel pain, and were originally thought to be its cause. Heel spurs occur at the origin of the flexor digitorum brevis muscle, which lies just beneath the plantar fascia. However, heel spurs are NOT the cause of heel pain associated with plantar fasciitis and for this reason, it is unnecessary for them to be removed. They probably form because of the local inflammation in the soft tissue, with the body forming bone mistakenly in its attempt to repair the plantar fascia.


NON-OPERATIVE MANAGEMENT

Initial management of plantar fasciitis involves a multi-modal non-operative approach. For 90% of people, this is all that is required. This initially includes analgesics, anti-inflammatories, massage, applying ice, and exercises that stretch the fascia and strengthen the muscles. Orthoses including cushioning heel pads and night splints may also be helpful, along with corticosteroid injections, and other alternative modalities such as ultrasound or shock wave therapy.

OPERATIVE MANAGEMENT

For those patients who have ongoing pain despite adequate non-operative management, operative intervention may be required. Conventionally this procedure has been performed through a large incision on the in-step of the foot. In recent times an endoscopic technique has been developed.

Our recent study of over 100 patients comparing the results of the traditional open procedure with the more modern endoscopic technique demonstrated the endoscopic procedure to be superior. This included improved patient satisfaction, less post-operative pain, quicker recovery times, and a lower complication rate.



The procedure involves making 2 x 10mm incisions; 1 on each side of the foot. This enables a telescopic camera and instrumentation to be inserted to complete the release of the plantar fascia. Patients are then allowed to weight bear as tolerated, and quite often have less pain than prior to the procedure within 1-2 weeks. By 4 weeks, patients are usually walking well, and may commence returning to sport by 6-12 weeks.

COMPLICATIONS

Complications are rare with this procedure, and in our series occurred in less than 5%. There is always a small risk of infection, nerve injury and blood clots, and anaesthetic problems with lower limb surgery and measures are taken to minimize these risks. A small proportion of patients may also experience a persistence of their pre-operative pain despite surgery. Finally, and uncommonly, discomfort in the outer part of the top of the foot may occur following any technique of plantar fascia release. Very seldom is this enough of a problem to require further surgery and generally improves with the use of insoles (orthotics). Overall, a successful outcome is achieved in approximately 90% of patients.

RECOVERY TIMES

Hospital stay	Day surgery
Rest & elevation	7 days
Crutches/frame	5-7 days
Foot swelling	4 weeks
Sport	6-12 weeks

TIME OFF WORK

Seated	1 week
Standing	3-4 weeks

This brochure is a brief overview of the surgical management of plantar fasciitis and not designed to be all-inclusive. If you have any further questions, please do not hesitate to contact your surgeon.

These notes have been prepared by orthopaedic surgeons at OrthoSport Victoria. They are general overviews and information aimed for use by their specific patients and reflects their views, opinions and recommendations. This does not constitute medical advice. The contents are provided for information and education purposes only and not for the purpose of rendering medical advice. Please seek the advice of your specific surgeon or other health care provider with any questions regarding medical conditions and treatment.

